



ELSEVIER

Contents lists available at ScienceDirect

Quaternary Science Reviews

journal homepage: www.elsevier.com/locate/quascirev

Climatic quantification and seasonality of the late MIS 3 in North China: A perspective from carbon and oxygen isotopes of fossil mammal teeth

Xu Wang^{a, b}, Jing Sun^c, Fred J. Longstaffe^d, Xuejun Gu^e, Shuisheng Du^{f, *}, Linlin Cui^{a, b}, Xiaozhong Huang^g, Zhongli Ding^{a, b}

^a Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, P.O. Box 9825, Beijing 100029, China

^b Innovation Academy for Earth Science, CAS, China

^c The Palace Museum, 4 Jingshan Qianjie, Beijing 100009, China

^d Department of Earth Sciences, The University of Western Ontario, London, Ontario N6A 5B7, Canada

^e Luoyang City Cultural Relics and Archaeology Research Institute, Luoyang 471000, China

^f School of History, Beijing Normal University, 19 Xijiekouwai Street, Beijing 100875, China

^g Key Laboratory of Western China's Environmental Systems (Ministry of Education) College of Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000, China

ARTICLE INFO

Article history:

Received 12 July 2021

Received in revised form

24 September 2021

Accepted 1 October 2021

Available online xxx

Handling Editor: Xiaoping Yang

Keywords:

Marine isotope stage 3

North China

Stable isotopes

Fossil mammal tooth

Climate

Seasonality

ABSTRACT

Marine Isotope Stage 3 (MIS 3, at 60–25 ka B.P.) was a short interstadial within the Last Glacial period. A generally warm and wet climatic condition during the MIS 3 period has been suggested by many previous studies. Quantitative climatic reconstruction and studies of seasonality, however, have been seldom attempted given the scarcity of useful proxy indicators. A full understanding of the climatic characteristics during this period is hampered by their absence. This is especially the case in the East Asian monsoon region in North China, which is very sensitive to climatic change. To help fill this knowledge gap, we have determined the stable carbon ($\delta^{13}\text{C}$) and oxygen isotope ($\delta^{18}\text{O}$) compositions of tooth enamel from fossil mammals (*Bubalus* sp [buffalo] and *Cervus elaphus* [red deer], dated at ~33–31 ka B.P.) recovered from Longquan Cave in North China. We use these data to constrain ecology, local air temperature and precipitation amounts during the late MIS 3. The average $\delta^{13}\text{C}$ (VPDB) of tooth enamel (−14.4‰ and −8.8‰, respectively for buffalo and red deer) indicate that C_3 plants comprised the main diet of animals living in this area. Pollen assemblages demonstrated that vegetation mainly consisted of grasses and herbs (~64%) with trees only accounting for about 26%. These results indicate a landscape of open steppe with sparsely distributed trees. The reconstructed climate derived from tooth enamel $\delta^{18}\text{O}$ was relatively colder and drier than present-day condition, with mean annual air temperature ranging from 4 to 13 °C and mean annual precipitation ranging from 360 to 670 mm. Moreover, the inferred summer and winter half-year temperatures were respectively 8–24 °C and −2 to 5 °C, which represents either similar or weaker seasonality than at the present time. The inferred cold and dry climate suggested by our study contrasts with the warm and wet conditions previously reconstructed using proxy indicators from bulk soil samples in North China. In comparison to the long-term, time-averaged climatic condition reflected by those soil proxies, the stable isotope compositions of fossil more likely recorded climatic conditions at annual or seasonal time-scale. The inferred annual cold/dry climate at Luanchuan Cave suggested by our data may indicate a short climatic cooling event from Greenland Interstadial (GIS) 5 to its following stadial. Mean annual temperature fluctuated by ~8 °C during this interstadial-stadial cycle.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

Marine Isotope Stage 3 (MIS 3), spanning 60–25 ka B.P., was a short interstadial stage within the Last Glacial period (Imbrie et al.,

* Corresponding author.

E-mail addresses: xuking@mail.iggcas.ac.cn (X. Wang), ssdu@bnu.edu.cn (S. Du).

1984). The climate during this stage was extremely unstable as reflected in the occurrence of millennial-scale climatic events, i.e., Dansgaard-Oeschger cycles and Heinrich stadials (Dansgaard et al., 1984, 1993; Heinrich, 1988; Johnsen et al., 1992; Stuiver and Grootes, 2000). Previous studies of MIS 3 have mainly focused on regional comparison of millennial-scale climatic variability and the mechanisms causing the abrupt climate events (Sánchez Goñi et al., 2002; Jiménez-Moreno et al., 2010; Sánchez Goñi and Harrison, 2010; Van Meerbeeck et al., 2011; Sionneau et al., 2013; Gibb et al., 2014). Quantitative reconstruction of regional climate (i.e., temperature and precipitation) and study of climate seasonality, however, have seldom been undertaken, in part due to the scarcity of useful proxy indicators or suitable materials to study. This is especially the case for the East Asian monsoon region in North China. This region is ideal for reconstructing paleoclimate and investigating the mechanisms of paleoclimatic evolution because it is more sensitive to abrupt climatic changes and seasonal variations (e.g., Wu et al., 2012).

The carbon ($\delta^{13}\text{C}$) and oxygen ($\delta^{18}\text{O}$) isotope compositions of tooth enamel (bioapatite) from mammals record information about the food and water consumed by animals (Longinelli, 1984; Bryant et al., 1994; Koch et al., 1998; Wang and Deng, 2005). Tooth enamel is very stable and hence resistant to carbon and oxygen isotope exchange after its formation, particularly during *post-mortem* and diagenetic processes (Quade et al., 1992). Hence, the isotopic information they recorded can directly reflect the climatic and ecological conditions in the study area during the animal's life, including vegetation type, temperature and precipitation (Wang et al., 1994). For example, the $\delta^{18}\text{O}$ of bioapatite from fossil tooth enamel has been used to quantitatively reconstruct air temperatures in the geological past (Delgado Huertas et al., 1997; Fricke and O'Neil, 1999; Tütken et al., 2007; Bernard et al., 2009; Amiot et al., 2010; Fabre et al., 2011; Amiot et al., 2011; Amiot et al., 2015). The $\delta^{13}\text{C}$ of structure carbonate from fossil tooth enamel has been employed to infer paleodiet and hence ecological information, i.e., the relative abundance of C_3/C_4 vegetation (e.g., Cerling et al., 1997; Koch, 1998; Deng et al., 2002). More recently, the $\delta^{13}\text{C}$ of fossil tooth enamel has also been employed to infer paleo-rainfall (Amiot et al., 2015).

In this study, we have determined the carbon and oxygen isotope compositions of bioapatite structural carbonate ($\delta^{13}\text{C}_\text{c}$ and $\delta^{18}\text{O}_\text{c}$, respectively) from tooth enamel along the growth bands of fossil teeth of buffalo (*Bubalus* sp.) and red deer (*Cervus elaphus*), excavated from the Longquan Cave Site, Luanchuan County, North China (Fig. 1a). These samples were buried *in situ* at ~33–31 ka B.P. during the late MIS 3 (Du et al., 2016). The oxygen isotope compositions of bioapatite phosphate ($\delta^{18}\text{O}_\text{p}$) were also measured and compared with the corresponding $\delta^{18}\text{O}_\text{c}$ to test for preservation of original oxygen isotope signals in the tooth enamel. We then use these data to quantitatively reconstruct temperature and rainfall, with special attention to climatic seasonality at the study site. These results provide insight into the MIS 3 climatic characteristics of North China at ~33–31 ka B.P.

2. Geological setting

Longquan Cave (33°47'24"N, 111°36'28"E) is located in Longquan Mountain Park in Luanchuan County, Luoyang City, Henan province (Fig. 1a). The climate of Luanchuan County is temperate and humid, fitting within the warm temperate continental monsoon climate classification. At present, the mean annual temperature (MAT) at the site is 12 °C, with a summertime (June–August) mean temperature of 22.9 °C and winter (December–February) mean temperature of 0.6 °C. The site's mean annual precipitation is 873 mm, and ~50% of the rainfall occurs in

summer, brought by the East Asian summer monsoon (Compilatory Commission of Annals of Luanchuan County, 1994).

Longquan Cave faces east and is situated about 6 m above the modern level of Yihe river. The cave is heavily damaged by erosion and by recent quarrying of its limestone roof and walls. Preserved layers were found at the back of the cave near the southern, western and northern walls. The site was divided into zones A–D and excavated within a grid of 1 m² sections (Fig. 1b and c). Zone C had the deepest stratigraphy, followed by zones B, A and D. Deposition in the cave can be subdivided into three stratigraphic layers, as described as below:

Layer 1 is uppermost, consisting of brownish-yellowish clay and brownish-yellowish silty clay, with a thickness of 0.3–1.05 m in zones A and C. In zone D, layer 1 was disturbed by modern human activity and its lower part is brownish-reddish clay with the thickness of only 0.05–0.15 m.

Layer 2 is hard, ~0.5–1.6 m thick and composed of reddish-brown clay, with a few pebbles in some areas. Most of the lithic artefacts and fossil animal bones recovered from the Longquan Cave were in this layer. The lithic assemblage (stone tools) belongs to the core-and-flake industry once prevalent in southern China (Du et al., 2016). The foragers who temporarily camped in the Longquan Cave used the simple core-and-flake technique to obtain flakes with sharp cutting edges. The faunal remains may originate from anthropogenic accumulations. A polished bone tool (i.e., a bone awl) was also discovered in layer 2 from zone D, which marks a significant innovation in tool-making technology. Two hearths were uncovered in layer 2 in zone A at depth of 100 mm and 200 mm, respectively. Collectively, the Longquan Cave provides evidence of traits regarded as similar to other modern human behavior, notably the distinct zoning of domestic areas for specific purposes, the use of hearths and rocks to store thermal energy, and the use of polished bone tools (Du et al., 2016).



Charcoal samples for AMS ¹⁴C dating were collected from the upper parts of layer 2 in zones A and D (Fig. 1b). A fossil bone collected from sub-layer 7 of layer 2 in zone D was also dated and had an age of ~33 ka cal BP. A total of eight radiocarbon dates were obtained. These radiocarbon dates were presented by Du et al. (2016), and showed that most of the depositional horizons span ~35 to 31 ka cal B.P. except for the lower hearth, which has a date of ~40 ka cal BP. It is therefore suggested initial human occupation occurred at ~40 ka cal BP, followed by a second period of activity at ~35 to 31 ka cal BP. Hence, the Longquan occupation falls within the MIS 3, a period that was generally marked by relatively rapid climatic fluctuations from colder to warmer on a millennial scale (e.g., Dansgaard et al., 1984, 1993). However, the climatic conditions that facilitated the Longquan occupation of the cave are not known. Since fossil teeth of buffalo and red deer were excavated from the lower part of layer 2 (i.e., sub-layers 5–7), they were assigned to an age of ~33–31 ka cal BP based on the two AMS ¹⁴C age control points in Zone D (Fig. 1b). Accordingly, the stable carbon and oxygen isotope compositions of enamel from these fossil tooth samples have been analyzed to help reconstruct the paleoclimate at this time.

Layer 3 is yellowish silt inter-bedded with brown-yellowish clay, and is only exposed in zone C (Fig. 1b). This deposit was excavated to ~0.7 m depth, but few lithics and fossil bones were found.

3. Materials and methods

3.1. Fossil teeth

Five fossil teeth of buffalo and two fossil teeth of red deer were analyzed in this study (Fig. S1). These teeth were visibly well-preserved. The teeth were all excavated from brown-reddish clay



in the layer 2, with six of them from sub-layers 5, 6 and 7 in zone D and one of them from sub-layer 5 in zone A (Table 1). Specifically, LCMD2(5):34 and LCMD2(6):19 are the third molar teeth (M3) from the upper jaws of two individual red deer collected from the sub-layer 5 and 6 of layer 2 in zone D, respectively. Tooth LCMA2(5):66–3 is the fourth deciduous premolar from the lower jaw of a buffalo, excavated from sub-layer 5 of layer 2 in zone A. Sample LCMD2(5):16 consists of three sequential molar teeth M1, M2 and M3 from the upper jaw of a buffalo unearthed from sub-layer 5 of the layer 2 in zone D. LCMD2(7):23–2 is the second molar tooth (M2) from the upper jaw of a buffalo, which was excavated from sub-layer 7 of layer 2 in zone D.

The samples described above belong to different tooth types, which would have had different formation times and growth rates. Intra-tooth isotopic analysis can therefore provide a wealth of information about seasonal changes over the period of tooth enamel development. For example, *C. elaphus* third molars (M3) erupt between 21 and 32 months, with enamel mineralization proceeded between 9 and 26 months (Godawa, 1989; Brown and Chapman, 1991; Azorit et al., 2002). Since deer calves are born in spring, the time sequence recorded in the M3 should therefore span early spring/summer/autumn/winter/late spring (Stevens et al., 2011).

The tooth formation sequence of *Bubalus* sp. is assumed to be similar to that of modern bison (Gadbury et al., 2000; Fricke and O'Neil, 1996). This sequence starts with growth of M1 *in utero* at the end of winter, followed by mineralization of M2 during the first year of the animal's life (summer to summer), then by M3, P2, P3 and P4 during the next year (Bernard et al., 2009). In this case, sample LCMD2(5):16, which contained M1, M2 and M3 in sequence, may represent a continuous record of the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ of diet and drinking water for a period of at least two years. The other samples (e.g., M2 and P4) probably documented one year's variation in isotopic compositions. We note, however, that the first molar (M1) forms, at least in part, *in vitro* from the body water of the mother. Hence, its $\delta^{18}\text{O}$ is unlikely to reflect the value of local meteoric water (Fricke and O'Neil, 1996). The second molar (M2) may start to form prior to weaning and its $\delta^{18}\text{O}$ may be slightly affected by isotopic composition of the milk the calf consumed during nursing (Gadbury et al., 2000).

3.2. Sampling

The teeth were first cleaned using a brush and distilled water to remove visible soil, and then air-dried in cool place. As needed, secondary deposits adhered to tooth surfaces were removed gently using dental drill with a 0.3 mm tip. After these cleaning procedures, ~10–15 mg of powder was sequentially drilled from the enamel using a diamond-incrusted drill bit. This subsampling was performed at ~2 mm intervals along the tooth growth axis from the apex to the cervix of the crown. A chronological record of isotopic composition during tooth mineralization can be obtained using this approach. The sampling groove for each subsample was milled perpendicular to the tooth length axis and encompassed the full enamel thickness. Care was taken to avoid dentin during sampling.

Previous studies have shown that amelogenesis (tooth enamel formation) involves a two-step process, which begins with the rapid deposition of an organic-rich, mineral-poor matrix and is then followed by a phase of progressive maturation that produces the fully mineralized enamel (Suga et al., 1970, 1987; Sakae and Hirai, 1982; Moss-Salentijn et al., 1997). During the maturation

stage, three mineralization fronts advance in various directions with some of them sub-parallel to the enamel-dentine junction (Suga, 1979, 1982). Each of the enamel subsamples collected here, and their carbon and oxygen isotope compositions therefore represent time-averaged signals of the body during the period of enamel mineralization at the subsample location (Zazzo et al., 2005). The pattern of isotopic variation encoded in tooth enamel, however, is attenuated compared to the true pattern of isotopic variation experienced by the animal (Passey and Cerling, 2002). Hence, it is not known exactly how much time is represented by each horizontal tooth subsample.

This dampening of the seasonal isotopic signal in tooth enamel hampers the ability to accurately reconstruct seasonal temperatures. Many efforts have been made to constrain the length of the enamel maturation process and correct for the dampening effect to recover a primary isotopic time-series (Passey and Cerling, 2002; Kohn, 2004; Zazzo et al. 2005, 2010; Bendrey et al., 2015; Traylor and Kohn, 2016). Those efforts provide important guidance for the paleoclimate reconstructions made using isotopic data gathered in the present study.

3.3. Chemical pretreatment and stable isotope determinations

Labile organic matter and secondary/adsorbed carbonates contained in tooth enamel can interfere with accurate measurement of $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$. Several pretreatment methods have been developed to remove these compounds before isotopic determination, including chemical oxidation using either H_2O_2 or NaOCl followed by acetic acid (of varying molarities) (Koch et al., 1997; Deng and Li, 2005; Wang and Deng, 2005; Snoeck and Pellegrini, 2015; Pellegrini and Snoeck, 2016). In our study, ~10 mg of enamel powder was pre-treated following the procedure of Koch et al. (1997), which uses very dilute chemical reagents to inhibit potential for enamel recrystallization in pretreatment solutions. Briefly, powders were washed with a 2% NaOCl solution at room temperature to remove organic matter, followed by a 0.1 M acetic acid solution to remove secondary carbonate. The powder/solution ratio was kept constant to 0.04 g/mL for both treatments. These treatments remove organic matter and secondary/adsorbed carbonates without substantial partial dissolution of hydroxylapatite (Koch et al., 1997). Each treatment lasted for 24 h, after which the subsamples were rinsed five times with distilled water and then dried at 50 °C.

About 2 mg aliquot of each pre-treated powder was analyzed using a GasBench II coupled with MAT 253™ isotope ratio mass spectrometer (Thermo Scientific™) at Laboratory for Environmental Isotope Geochemistry, Institute of Geology and Geophysics, CAS (LEIG-IGGCAS), following a procedure adapted after Spötl and Vennemann (2003). Five drops of 100% phosphoric acid were added allowing the samples to react at 72 °C for 1 h under a He atmosphere before starting 10 measurement cycles of the isotopic composition of the produced CO_2 . The measured carbon and oxygen isotope compositions were normalized to VPDB scale using two reference materials as scale end-points, including one international reference material NBS-19 calcite ($\delta^{13}\text{C} = +1.95\text{‰}$ and $\delta^{18}\text{O} = -2.20\text{‰}$) and one national reference material of China GBW04416 calcite ($\delta^{13}\text{C} = -10.85\text{‰}$ and $\delta^{18}\text{O} = -12.40\text{‰}$). Another national reference material of China, GBW04405 calcium carbonate ($\delta^{13}\text{C} = +0.57\text{‰}$ and $\delta^{18}\text{O} = -8.49\text{‰}$), was inserted into the measurement sequence as a quality control. Reproducibility of $\delta^{13}\text{C}$

Fig. 1. (a) The location of Longquan Cave in Luanchuan County, Henan Province. (b) Cross-section of the different zones inside Longquan Cave. AMS ^{14}C age data were shown to constrain the time-scale of the horizons where fossil teeth were excavated. (c) Photograph of Longquan Cave (credit of S.S. Du). Showing the locations of zones A to D, as described in the text. The cave roof has collapsed.

Table 1
Mean stable carbon and oxygen isotope results for tooth enamel from red deer (*Cervus elaphus*) and buffalo (*Bubalus* sp.).

Sample No.	Horizon	Species	Tooth type	$\delta^{13}\text{C}_\text{C}$ (‰, VPDB)	$\delta^{18}\text{O}_\text{C}$ (‰, VSMOW)	Number of $\delta^{13}\text{C}_\text{C}/\delta^{18}\text{O}_\text{C}$ microsamples (n)	$\delta^{18}\text{O}_\text{P}$ (‰, VSMOW)	Number of $\delta^{18}\text{O}_\text{P}$ microsamples (n)	$\delta^{18}\text{O}_\text{C} - \delta^{18}\text{O}_\text{P}$ (‰)
LCMD2(5):34	2(5)	<i>Cervus elaphus</i>	^a UP M3	-13.2	+23.2	17	+13.9	8	9.3
LCMD2(6):19	2(6)	<i>Cervus elaphus</i>	UP M3	-12.9	+23.9	11	+16.1	5	7.8
LCMA2(5):56-3	2(5)	<i>Bubalus</i> sp.	LR dP4	-13.5	+23.5	14	+13.3	13	10.2
LCMD2(5):16	2(5)	<i>Bubalus</i> sp.	UP M1	-13.2	+20.4	9	+11.9	9	8.4
			UP M2	-13.1	+20.4	11	+12.9	13	7.5
			UP M3	-13.1	+19.3	9	+12.5	12	6.8
LCMD2(7):23-2	2(7)	<i>Bubalus</i> sp.	UP M2	-10.3	+22.4	34	+16.6	32	5.8

Each isotopic result is the calculated mean for whole-tooth enamel from several microsamples (n). Individual data are tabulated in [Supplementary Table S1](#).

^a LR and UP represent lower and upper jaw, respectively; M denotes molar tooth and dP denotes deciduous premolar.

and $\delta^{18}\text{O}_\text{C}$ for this standard was better than $\pm 0.1\%$ and $\pm 0.2\%$ (1SD, $n = 10$), respectively. We also performed duplicate analyses of ten enamel sub-samples to determine their isotopic reproducibility. The maximum difference between duplicate enamel analyses was 0.2% for $\delta^{13}\text{C}$ and 0.25% for $\delta^{18}\text{O}$. The oxygen isotope compositions were then converted from the VPDB scale to the VSMOW scale using the equation: $\delta^{18}\text{O}_\text{VSMOW} = 1.03091 \times \delta^{18}\text{O}_\text{VPDB} + 30.91$ (Coplen et al., 1983).

Extraction of bioapatite phosphate consists of isolating the phosphate anionic complexes using acid dissolution and anion-exchange resin and its precipitation as silver phosphate, according to a protocol reported by Crowson et al. (1991) and slightly modified by Lécuyer et al. (1993). Silver phosphate was quantitatively precipitated in a thermostatic bath set at 70°C . The silver phosphate was then recovered by filtration, washing with double-deionised water, and then drying at 50°C . Extraction of bioapatite phosphate from one buffalo tooth (LCMD2(7):23–2) was carried out at LEIG-IGGCAS and its $\delta^{18}\text{O}_\text{P}$ measured using a high temperature conversion elemental analyzer (TC/EA) linked to MAT253 isotope ratio mass spectrometer (Thermo Scientific™). Extraction and isotopic analysis of bioapatite phosphate from the other six fossil teeth were performed at the Laboratory for Stable Isotope Science, The University of Western Ontario, Canada (LSIS UWO). The $\delta^{18}\text{O}_\text{P}$ of those silver phosphate subsamples were measured using a TC/EA coupled with a Delta™ Plus XL isotope ratio mass spectrometer (Thermo Scientific™).

Values of $\delta^{18}\text{O}_\text{P}$ are reported relative to VSMOW, as calibrated using IAEA-CH-6 ($\delta^{18}\text{O} = +36.4\%$; Flanagan and Farquhar, 2014) and Aldrich Silver Phosphate–98%, Batch 03610 EH ($\delta^{18}\text{O} = +11.2\%$; Webb et al., 2014). Silver phosphate precipitated from standard NBS120c (Miocene phosphorite from Florida: $\delta^{18}\text{O}_\text{P} = +21.7$; Amiot et al., 2011) was analyzed along with the silver phosphate samples derived from the fossil teeth as a check on the calibration. The reproducibility of $\delta^{18}\text{O}$ results for standards (Aldrich, $n = 18$; IAEA-CH-6, $n = 10$; NBS120c, $n = 5$) and duplicate sample pairs ($n = 18$) was better than $\pm 0.3\%$ (SD) in all cases.

3.4. Pollen analysis

A total of eight soil samples were collected from Longquan Cave section for pollen analysis. The samples were processed using standard methods, including treatment with HCl (10%), hot KOH (10%) for 5–10 min, HF (64%), and fine sieving in an ultrasonic bath (Faegri et al., 1989). At the beginning of the procedure one tablet of *Lycopodium* spores was added to each sample to enable calculation of the pollen concentration (Maher, 1981). The pollen concentrates were mounted in glycerol and examined at $\times 400$ magnification

using a Nikon optical microscope (Model: ECLIPSE-80i). Where appropriate, magnifications of $\times 600$ and/or $\times 1000$ were used to examine pollen surface sculpturing. Identification was aided by reference to Wang et al. (1995), together with comparisons with modern pollen reference collections. Pollen counts consisted of ~ 200 – 400 terrestrial pollen grains except for two samples at lower part of the sequence where the pollen abundance was much lower. Pollen percentages were calculated based on the sum of all terrestrial pollen taxa present.

4. Results

All stable isotope data ($\delta^{13}\text{C}_\text{C}$, $\delta^{18}\text{O}_\text{C}$ and $\delta^{18}\text{O}_\text{P}$) for each tooth are listed in [Table S1](#). [Fig. 2](#) illustrates changes in $\delta^{13}\text{C}_\text{C}$ (VPDB) and $\delta^{18}\text{O}_\text{C}$ (VSMOW) along the growth axis of each tooth. For the two fossil teeth of red deer, $\delta^{13}\text{C}_\text{C}$ ranges from -13.5 to -12.4% and $\delta^{18}\text{O}_\text{C}$ ranges from $+22.2$ to $+24.5\%$. Neither tooth for the red deer recorded a full range of seasonal changes in isotopic composition, possibly because the teeth were relatively short (< 20 mm) and the crowns were partially worn.

Larger variations are observed for $\delta^{13}\text{C}_\text{C}$ (-14.3 to -8.8%) and $\delta^{18}\text{O}_\text{C}$ ($+17.6$ to $+25.1\%$) of the five fossil buffalo teeth. These variations can be attributed to seasonal changes. Tooth LCMA2(5):56–3 is a deciduous premolar and documented a half cycle of seasonal changes in its $\delta^{18}\text{O}_\text{C}$ and $\delta^{13}\text{C}_\text{C}$. By comparison, tooth LCMD2(7):23–2 measures ~ 70 mm from cervix to apex and its $\delta^{18}\text{O}_\text{C}$ exhibits a sinusoidal pattern of variation attributed to a full seasonal cycle. By contrast, the $\delta^{13}\text{C}_\text{C}$ of this tooth showed more frequent fluctuations about -10.3% . Sample LCMD2(5):16 contained 3 M teeth from the first to the third molar (designated as LCMD2(5):16 M1, LCMD2(5):16 M2 and LCMD2(5):16 M3, respectively). Their isotopic data can be combined to generate a composite isotopic seasonal time series. Because the M1 tooth of cattle is known to form prior to birth (Fricke and O'Neil, 1996), it must have formed *in vitro* from the body water of mother. We assume that the same tooth eruption schedule holds for its close relative, the buffalo. Therefore, its M1 tooth can be used to investigate seasonality during the birthing period, followed by the M2 and M3 whose isotopic data provide a good illustration of seasonal variations ([Fig. 2](#)).

[Fig. 3](#) compares intra-tooth variation in $\delta^{18}\text{O}_\text{P}$ and $\delta^{18}\text{O}_\text{C}$ for each fossil tooth. Overall, $\delta^{18}\text{O}_\text{P}$ generally describes the same pattern as $\delta^{18}\text{O}_\text{C}$. The $\delta^{18}\text{O}_\text{P}$ for the two red deer teeth range from $+12.6$ to $+14.4\%$ for LCMD2(5):34 and from $+13.9$ to $+19.3\%$ for LCMD2(6):19. The latter tooth displays a sharp decrease in $\delta^{18}\text{O}_\text{P}$ from the apex towards the cervix, a pattern also observed for $\delta^{18}\text{O}_\text{C}$.

The $\delta^{18}\text{O}_\text{P}$ of the LCMA2(5):56–3 pre-molar buffalo tooth (dP4)




Fig. 2. Intra-tooth variations in stable carbon and oxygen isotope compositions of bioapatite structural carbonate ($\delta^{13}\text{C}$ in ‰, VPDB; $\delta^{18}\text{O}$ in ‰, VSMOW) in tooth enamel of seven fossil teeth versus distance (in mm) from the cervix.

fluctuates widely (+11.9 to +14.8‰), while increasing overall with decreasing distance from the cervix. The $\delta^{18}\text{O}_\text{P}$ for the long buffalo tooth LCMD2(7):23-2 shows a much more restricted range (+15.7 to +17.2‰; mean = +16.6‰). Its narrow range of $\delta^{18}\text{O}_\text{P}$ nevertheless registers a full seasonal cycle of variation. The molar (M1-M3) teeth set of LCMD2(5):16 show a larger oscillation in $\delta^{18}\text{O}_\text{P}$ from +10.6 to +17.9‰, which is suggestive of seasonal variation. The M1 has very low $\delta^{18}\text{O}_\text{P}$ showing little variation (+11.3 to +12.5‰), the M2 shows the largest variation in $\delta^{18}\text{O}_\text{P}$ (+10.6 to +17.9‰), rising to maximum values near its center, followed by a large decrease, and the M3 exhibits a gradual decrease from +13.6 to +10.7‰.

Table 1 lists mean values of $\delta^{13}\text{C}_\text{C}$, $\delta^{18}\text{O}_\text{C}$ and $\delta^{18}\text{O}_\text{P}$ for each tooth. The mean $\delta^{13}\text{C}_\text{C}$ varies from -13.5 to -12.9‰, except one more positive value of -10.3‰ (buffalo tooth LCMD2(7):23-2). The mean $\delta^{18}\text{O}_\text{C}$ varies from +19.3 to +23.9‰. The mean $\delta^{18}\text{O}_\text{P}$ ranges from +11.9 to +16.6‰. The mean value of $\delta^{18}\text{O}_\text{C}$ - $\delta^{18}\text{O}_\text{P}$ varies from +5.8 to +10.2‰, with no apparent inter-species differences.

The pollen samples were composed of 22 plant species, mainly trees, shrubs, grasses and herbs, and a few ferns (Fig. 4). Trees

consisted of spruce (*Picea*), birch (*Betula*), elm (*Ulmus*), willow (*Salix*), cypress (*Cupressaceae*) and a small quantity of poplar/aspens/cottonwood (*Populus*) and comprise ~27% of the pollen count. Shrubs included *Elaeagnaceae*, *Ephedra* and *Rosaceae*, but account for only ~3% of the pollen count. Of the grasses and herbs, *Ranunculaceae* and *Gramineae* most frequently occurred, with sporadic emergence of *Chenopodiaceae*, *Artemisia*, *Cypereceae*, *Compositae*, *Leguminosae*, and *Typha*. The grasses and herbs comprise ~64% of the pollen count. Spores make up ~6% of the pollen count.

5. Discussion

5.1. Factors influencing stable carbon and oxygen isotope compositions of mammalian tooth enamel

For herbivores like red deer and buffalo, their tooth enamel $\delta^{13}\text{C}_\text{C}$ is generally governed by the carbon isotope compositions of their plant forage. Most modern plants photosynthesize using either the C_3 pathway (C_3 plants) or the C_4 pathway (C_4 plants). Modern C_3 plants, which include trees, most shrubs and cool season




Fig. 3. Intra-tooth variations in oxygen isotope compositions of bioapatite phosphate ($\delta^{18}\text{O}_p$ in ‰, VSMOW), compared to data for bioapatite structural carbonate ($\delta^{18}\text{O}_c$ in ‰, VSMOW) in tooth enamel of seven fossil teeth. The $\delta^{18}\text{O}_p$ and $\delta^{18}\text{O}_c$ show generally similar changing patterns in the same tooth. Seasons were assigned according to the peaks and troughs of $\delta^{18}\text{O}_p$ and $\delta^{18}\text{O}_c$ curves, as shown by horizontal bars.

grasses, have $\delta^{13}\text{C}$ ranging from about -35 to -20 ‰, with an average of -27 ‰ (O'Leary, 1981; O'Leary, 1988; Farquhar et al., 1989). By contrast, modern C_4 plants, which are mostly warm season grasses, have $\delta^{13}\text{C}$ ranging from about -17 to -9 ‰, averaging -13 ‰. These isotopic signals are passed along the food chain to animal tissues with further isotopic fractionation. The average carbon isotope fractionation factor between bioapatite structural carbonate and diet is ~ 14 ‰ for ungulate mammals (Cerling and Harris, 1999; Passey et al., 2005). Therefore, a $\delta^{13}\text{C}_c$ of -13 ‰ would ideally indicate a dietary intake of modern pure C_3 plants whereas a $\delta^{13}\text{C}_c$ of $+1$ ‰ would indicate a pure diet of modern C_4 vegetation (e.g., Wang et al., 2008). This provides a guideline for inferring the proportions of C_3 and C_4 plants in a herbivore's diet, as has been done in many previous studies (e.g., MacFadden et al., 1996; Cerling et al., 1997; Wang and Deng, 2005). Other factors, however, such as soil moisture, aridity and atmospheric $p\text{CO}_2$ can induce a large variability (up to several per mil) in the $\delta^{13}\text{C}$ of plants (e.g., Farquhar et al., 1989; Cerling et al., 1997),

which need to be considered when inferring the C_3/C_4 from $\delta^{13}\text{C}_c$. In addition, because of the Suess Effect (e.g., Keeling, 1979; Marino et al., 1992), the ranges and averages of vegetation $\delta^{13}\text{C}$, and therefore $\delta^{13}\text{C}_c$, would have been higher by ~ 1.5 – 2 ‰ during MIS 3, i.e., -11.5 ‰ and $+2.5$ ‰ for teeth of mammals ingesting pure C_3 and C_4 plants, respectively.

The oxygen isotope composition of tooth enamel in mammals is directly linked to the $\delta^{18}\text{O}$ of body water because mammalian tooth bioapatite is formed at constant body temperature (Longinelli, 1984; Luz et al., 1984; lacumin et al., 1996). Body water $\delta^{18}\text{O}$ is controlled by several factors, including the $\delta^{18}\text{O}$ of drinking water and water in food (i.e., plants for herbivores), physiological processes and dietary or drinking behavior (Longinelli, 1984; Luz et al., 1984; Kohn et al., 1996). Several studies have shown that the $\delta^{18}\text{O}$ signal in tooth enamel from large animals (>1 kg) is dominated by meteoric water (mw) because most ingested water comes from that source (Longinelli, 1984; Luz et al., 1984; Bryant et al., 1996). Because $\delta^{18}\text{O}_{\text{mw}}$ correlates with climatic variables including




Fig. 4. Pollen percentages in bulk samples from the second layer of Longquan Cave deposits.

temperature, seasonality of rainfall and rainfall amount (Dansgaard, 1964; Rozanski et al., 1993), the $\delta^{18}\text{O}$ of enamel can be used to reconstruct paleoclimatic conditions during tooth growth (e.g., Longinelli, 1984; Bryant et al., 1994; Bryant et al., 1996; Wang et al., 2008). The most common application of this $\delta^{18}\text{O}_\text{C}$ and $\delta^{18}\text{O}_\text{P}$ proxy has been to infer local surface temperatures in the geological past (e.g., Delgado Huertas et al., 1997; Fricke and O'Neil, 1999; Wang and Deng, 2005; Tütken et al., 2007; Bernard et al., 2009; Amiot et al., 2010; Fabre et al., 2011; Amiot et al., 2011; Amiot et al., 2015).

Another significant use of tooth enamel $\delta^{13}\text{C}_\text{C}$ and $\delta^{18}\text{O}_\text{C}/\delta^{18}\text{O}_\text{P}$ has been to trace seasonal variations in diet and climate during tooth growth through analysis of serial enamel samples from an individual tooth (e.g., Fricke and O'Neil, 1996; Dettman et al., 2001; Bernard et al., 2009; van Dam and Reichart, 2009; Metcalfe et al., 2011). This approach is undertaken because mammalian tooth enamel forms incrementally from the crown to the cervix of the tooth, recording a series of $\delta^{13}\text{C}_\text{C}$ and $\delta^{18}\text{O}_\text{C}/\delta^{18}\text{O}_\text{P}$ over time. These isotopic compositions may vary by several per mil along the growth axis within a single tooth and thus provide an excellent record of dietary and/or climatic seasonality, albeit that the isotopic signal might be attenuated due to the time-averaging effect induced by the sampling technique (Passey and Cerling, 2002), as discussed earlier. Migratory behavior can also make seasonal inferences from such isotopic data more complicated. Red deer, however, are not a migratory species (Fabre et al., 2011) and their tooth $\delta^{18}\text{O}$ has been shown to record perfect seasonal variation (Stevens et al., 2011). In contrast, the migratory behavior of the Chinese buffalo is not fully known. Because buffalo is water-dependent animal, it relies mostly on standing water sources (e.g., lakes). The Yihe River, which is located nearby our study area, likely met this demand, thus minimizing demands on the buffalo to travel long distances. In addition, the aquatic plants associated with the river system may have offered preferred forage to the buffalo.

5.2. Carbon isotopes, paleodiet and paleoecology at the study site

The intra-tooth $\delta^{13}\text{C}_\text{C}$ of the red deer varied only slightly (-13.5 to -12.4‰) (Fig. 2 and Table S1), which are more negative than the MIS 3 end-member value of -11.5‰ for a dietary intake of pure C_3

plants. In contrast, much larger variation in intra-tooth $\delta^{13}\text{C}_\text{C}$ (-14.3 to -8.8‰) was observed for buffalo. The less negative values of $\delta^{13}\text{C}_\text{C}$ (-11.2 to -8.8‰) were obtained only for the M2 tooth in sample LCMD2(7):23–2. The range in $\delta^{13}\text{C}_\text{C}$ (-14.3 to -12.5‰) for other buffalo teeth is similar to that determined for the red deer. The higher $\delta^{13}\text{C}_\text{C}$ of tooth LCMD2(7):23–2 could have been caused by intake of C_4 plants. However, if this was the reason, the higher values of $\delta^{13}\text{C}_\text{C}$ would be expected during summer (e.g., period of peaks in the intra-tooth $\delta^{18}\text{O}_\text{C}$ curve) when C_4 plants normally reach peak abundance (Yang et al., 2012). That appears not to have been the case, however, because the higher $\delta^{13}\text{C}_\text{C}$ values corresponded to a trough in $\delta^{18}\text{O}_\text{C}$ curve (Fig. 2). Instead, we consider that the shift in $\delta^{13}\text{C}_\text{C}$ towards higher values in this tooth enamel probably resulted from relatively dry conditions during this period of tooth formation.

While a general bioapatite-diet ^{13}C enrichment of $\sim 14\text{‰}$ is generally assumed for ungulate mammals, some taxa-specific enrichment factors have been reported. Cerling and Harris (1999), for example, reported a value of 14.8‰ for buffalo, similar to the value of 14.6‰ reported for cattle (*B. taurus*) by Passey et al. (2005). Using this enrichment factor, the inferred dietary $\delta^{13}\text{C}$ for the buffalo ranged from -29.1 to -23.6‰ (i.e., $\delta^{13}\text{C}_{\text{diet-of-bufflo}} = \delta^{13}\text{C}_\text{C} - 14.8$). In the absence of a specific enrichment factor for red deer, the value for cattle has been used here, as both are ruminants, and may have had similar ecological habits. The resulting calculated dietary $\delta^{13}\text{C}$ for the red deer varies from -28.1 to -27.0‰ (i.e., $\delta^{13}\text{C}_{\text{diet-of-red deer}} = \delta^{13}\text{C}_\text{C} - 14.6$). These calculations suggest that these herbivores fed predominantly on C_3 plants during the late MIS 3. Such pure or nearly pure C_3 diets for these two different herbivores suggest that the ecosystems surrounding the study site consisted primarily of C_3 plants at $\sim 33\text{--}31$ ka B.P.

At present, our study area lies in the maximum C_4 biomass zone, i.e., the region from 31° to 40°N in China (e.g., Rao et al., 2008; Wang et al., 2000). However, the study site is located in the Funiu Mountain area (at the eastern part of Qingling Mountains) at an average elevation of ~ 800 m. Local climate at such a high elevation may be more suitable for the growth of C_3 plants. Therefore, the inferred C_3 vegetation in this study may represent the local landscape inhabited by the buffalo and red deer. It cannot be used to reflect a general feature of C_3/C_4 vegetation in northern China. A C_3

vegetation-dominated environment in this region could vary from dense forest to steppe with sparse trees. The range of carbon isotope compositions calculated here (range: -29.1 to -23.6‰) is more typical of a steppe environment. Dense forest vegetation typically has more negative $\delta^{13}\text{C}$ (as low as -34‰) due to the influence of soil respiration associated with the canopy effect (Schleser and Jayasekera, 1985; Sternberg et al., 1989; Van der Merwe and Medin, 1989; Drucker et al., 2008; Bonafini et al., 2013).

The possibility of a steppe-like environment is confirmed by the pollen data for our study site (Fig. 4). Grasses and herbs comprised the majority ($\sim 64\%$) of the pollen count, whereas trees and shrubs were in a distinct minority ($\sim 30\%$). Collectively, the pollen data indicate an open steppe landscape with a sparse distribution of trees, suggestive of a relatively cold and dry climate (e.g., Jiang and Ding, 2005). The inferred landscape is in accordance with red deer's preferred ecological niche, red deer like a boreal forest-meadow habitat, including alpine forest, shrub and grassland environments (Zhao, 1974).

5.3. Climate reconstruction and seasonality in North China during the late MIS 3

5.3.1. Tooth enamel $\delta^{13}\text{C}_\text{C}$ and paleorainfall reconstruction

Water availability is an important factor that influences plant carbon isotope discrimination in plants (Deines, 1980; Francey and Farquhar, 1982). A restricted soil water supply causes plants to close their stomata to conserve water for photosynthesis. This process in turn reduces the intercellular partial pressure of CO_2 , which causes plant $\delta^{13}\text{C}$ to rise accordingly. Recent investigations of modern ecosystems in northern China have shown that plant $\delta^{13}\text{C}$ shifts toward more negative values with an increase in mean annual precipitation (MAP) (Wang et al., 2003, 2013a; Liu et al., 2005). Wang et al. (2003), for example, observed a 0.47‰ negative shift in $\delta^{13}\text{C}$ of C_3 plants with an annual 100 mm increase in rainfall. Several recent studies have used such correlations to infer precipitation amount from the $\delta^{13}\text{C}$ of C_3 plants (e.g., Wang et al., 2013a; Amiot et al., 2015).

We initially attempted to adopt a previously established relationship between local MAP and the carbon isotope composition of modern C_3 plants in North China (Wang et al., 2013a) to estimate local precipitations at our study site. However, a recent study has shown that there is a strong positive correlation between the $\delta^{13}\text{C}$ of C_3 plants and mean annual temperature in north China (0.104‰ per $^\circ\text{C}$; Wang et al., 2013b). We have corrected for this effect, after removal of two outliers from the Wang et al. (2013a) dataset, to obtain the following regression (Fig. 5):

$$\text{MAP} = 90.862(\pm 14.050) \times \Delta_{\text{leaf}} - 1360.154(\pm 270.574) \quad (R^2 = 0.59) \quad (1)$$

where $\Delta_{\text{leaf}} = (\delta^{13}\text{C}_{\text{atm}} - \delta^{13}\text{C}_{\text{leaf}})/(1 + \delta^{13}\text{C}_{\text{leaf}}/10^3)$, $\delta^{13}\text{C}_{\text{atm}}$ is $\delta^{13}\text{C}$ of atmospheric CO_2 , and $\delta^{13}\text{C}_{\text{leaf}}$ represents the C_3 plant $\delta^{13}\text{C}$ corrected for the temperature effect.

Average local precipitation amounts calculated using equation (1) are listed in Table 2, using $\delta^{13}\text{C}_{\text{atm}} = -6.6\text{‰}$, as estimated for the late MIS 3 bases on CO_2 trapped in EPICA Dome C ice core (Lourantou et al., 2010), and the calculated mean $\delta^{13}\text{C}$ of fossil tooth enamel. All calculated values for our study site are below 670 ± 169 mm, with most of them ranging between 590 and 670 mm. Despite that the first molar (M1) of buffalo sample LCMD2(5):16 was formed *in utero* and the second molar (M2) was partly subject to nursing effects, their $\delta^{13}\text{C}$ generate MAP of 639 ± 161 mm and 627 ± 158 mm, respectively, very close to that obtained from the third molar (M3: 633 ± 160 mm). Moreover, the $\delta^{13}\text{C}$ for teeth of both red deer and buffalo from the sub-layer 5 yield a consistent MAP of ~ 630 mm (Table 2). A comparable MAP

(596 ± 150 mm) was also determined for red deer tooth (LCMD2(6):19) collected in sub-layer 6. However, the MAP inferred from buffalo tooth LCMD2(7):23–2 collected in the sub-layer 7 was substantially lower (363 ± 91 mm). This may reveal highly variable fluctuations in precipitation amounts at different times during late MIS 3. Collectively, inferred MAP are lower than the average annual amount of modern precipitation (873 mm). This relatively dry condition is consistent with the development of open steppe in the region, as also suggested by the pollen assemblages.

5.3.2. Tooth enamel $\delta^{18}\text{O}_\text{P}$ and paleotemperature reconstruction

As discussed earlier, $\delta^{18}\text{O}_\text{P}$ and $\delta^{18}\text{O}_\text{C}$ of mammal teeth are ultimately controlled by $\delta^{18}\text{O}_{\text{mw}}$ (e.g., Cormie et al., 1994; Kohn et al., 1996; Kohn and Cerling, 2002; Straight et al., 2004). In the current study, $\delta^{18}\text{O}_{\text{mw}}$ has been estimated from the $\delta^{18}\text{O}_\text{P}$ using various phosphate oxygen isotope-water geothermometers. In the absence of a buffalo-specific equation, the bison bioapatite enamel-water geothermometer of Bernard et al. (2009), following Hoppe (2006), was used, recognizing that buffalo and bison have similar body size and physiology:

$$\delta^{18}\text{O}_{\text{mw}} = 1.18(\pm 0.07) \times \delta^{18}\text{O}_\text{P} - 27.2(\pm 1.1) \quad (R^2 = 0.8) \quad (2)$$

An oxygen isotope bioapatite enamel-water geothermometer for red deer was developed based on the data of D'Angela and Longinelli (1990):

$$\delta^{18}\text{O}_{\text{mw}} = 0.879(\pm 0.033) \times \delta^{18}\text{O}_\text{P} - 22.50(\pm 0.52) \quad (R^2 = 0.99) \quad (3)$$

Amiot et al. (2004) developed a transfer function of $\delta^{18}\text{O}_{\text{mw}}$ to mean annual air temperature (MAT) by analyzing the monthly averages of temperature and isotopic data provided by the global network of stations operated by the IAEA–WMO (International Atomic Energy Agency–World Meteorological Organization):

$$\delta^{18}\text{O}_{\text{mw}} = 0.49(\pm 0.03) \times \text{MAT} - 14.18(\pm 0.52) \quad (R^2 = 0.81) \quad (4)$$

Using equations (2)/(3) and (4), the calculated MAT at our study site ranged from 2.2 ± 0.9 $^\circ\text{C}$ to 13.3 ± 1.2 $^\circ\text{C}$ during this portion of MIS 3 (Table 2). The lowest MAT value, however, was calculated for the $\delta^{18}\text{O}_\text{P}$ of the M1 tooth, which was influenced by the mother body fluid and therefore is disregarded. The remaining range (3.4 ± 0.9 to 13.3 ± 1.2 $^\circ\text{C}$) still represents a very large inter-annual variation in MAT. While the errors overlap, the mean MAT of 7.7 ± 4.0 (1SD) $^\circ\text{C}$ is likely diagnostic of temperatures lower than the present-day annual value of ~ 12 $^\circ\text{C}$.

The seasonality of air temperatures (T_{air}) at Longquan Cave site during the late MIS 3 can perhaps be better estimated using linear regression equations of $\delta^{18}\text{O}_{\text{mw}}-T_{\text{air}}$ derived from data obtained closer to the study area rather than from average values inferred from global datasets. We calculated such annual and seasonal regressions using a sub-dataset (1985–2003) for stations in China available through the IAEA-GNIP/ISOHIS dataset (World Data Center for Paleoclimatology, 2006) (Fig. 6a). Four stations were excluded, either due to a strong, year-round amount effect (Guangzhou, Haikou and Hongkong) or altitude effect (Lasha). Their exclusion improved the $\delta^{18}\text{O}_{\text{mw}}-T$ correlation.

The annual and winter regression equations obtained were:

Annual:

$$\delta^{18}\text{O}_{\text{mw}} = 4.765(\pm 0.548) \times \ln(\text{MAT}) - 19.578(\pm 1.362) \quad (R^2 = 0.75) \quad (5)$$

Winter half-year (October–March):

$$\delta^{18}\text{O}_{\text{mw}} = 0.681(\pm 0.052)T_{\text{air}} - 12.689(\pm 0.407) \quad (R^2 = 0.87) \quad (6)$$




Fig. 5. Correlation between Mean Annual Precipitation (MAP) (mm) and the carbon isotope fractionation (‰) between modern C_3 plants and atmospheric carbon dioxide ($\Delta_{\text{leaf-atm CO}_2}$). Data are from Wang et al. (2003).

Table 2

Inferred $\delta^{13}\text{C}_{\text{leaf}}$, $\Delta_{\text{leaf-atm CO}_2}$, mean annual precipitation (MAP), $\delta^{18}\text{O}_{\text{mw}}$ and mean annual temperature (MAT) based on $\delta^{13}\text{C}_C$ and $\delta^{18}\text{O}_P$ of red deer and buffalo.

Sample No.	Mammal species	Mean $\delta^{13}\text{C}_C$ (‰, VPDB)	$\delta^{13}\text{C}_{\text{leaf}}$ (‰, VPDB)	$\Delta_{\text{leaf-atm CO}_2}$ (‰)	MAP (eq. (1)) (mm)	Mean $\delta^{18}\text{O}_P$ (‰, VSMOW)	$\delta^{18}\text{O}_{\text{mw}}$ (‰, VSMOW)	MAT (eq. (4)) (°C)	MAT (eq. (5)) (°C)
LCMD2(5):34	Red deer	-13.2	-27.8	21.8	620 ± 156	+13.9	-10.3	7.9 ± 1.0	7.5 ± 2.5
LCMD2(6):19	Red deer	-12.9	-27.5	21.5	596 ± 150	+16.1	-8.4	11.8 ± 1.1	11.3 ± 4.1
LCMA2(5):56-3	Buffalo	-13.5	-28.3	22.3	670 ± 169	+13.3	-11.5	5.5 ± 1.0	5.8 ± 1.8
LCMD2(5):16M1	Buffalo	-13.2	-28.0	22.0	639 ± 161	+11.9	-13.1	2.2 ± 0.9	4.1 ± 1.2
LCMD2(5):16M2	Buffalo	-13.1	-27.9	21.9	627 ± 158	+12.9	-12.0	4.4 ± 1.0	5.2 ± 1.6
LCMD2(5):16M3	Buffalo	-13.1	-27.9	21.9	633 ± 160	+12.5	-12.5	3.4 ± 0.9	4.7 ± 1.4
LCMD2(7):23-2	Buffalo	-10.3	-25.1	19.0	363 ± 91	+16.6	-7.7	13.3 ± 1.2	13.1 ± 4.9

There was no significant correlation of $\delta^{18}\text{O}_{\text{mw}}$ with summer T_{air} , likely because of an apparent amount effect that produced a lowering of $\delta^{18}\text{O}_{\text{mw}}$ in the summer period. This effect is illustrated in Fig. 6b for Zhengzhou City, which is located close to the study area. Summer temperature (half year: April to September) was therefore calculated from using the inferred winter half-year T_{air} and local MAT. We also reconstructed MAT using equation (5), which produced results very close (maximum difference of ~ 1.9 °C) to those obtained using the global dataset (equation (4)) (Table 2).

Despite amount effect exists, $\delta^{18}\text{O}_{\text{mw}}$ in the early and late summer (except for the middle part) is still higher than that in winter. In this case, summer and winter seasons of enamel formation were assigned according to peaks and troughs, respectively, in the intra-tooth $\delta^{18}\text{O}_P$ and $\delta^{18}\text{O}_C$ curves (Fig. 3). As mentioned earlier, however, the dentition signal of seasonal variation in

$\delta^{18}\text{O}_{\text{mw}}$ is both damped and delayed due to contributions of atmospheric oxygen, residence time of oxygen in the body, enamel mineralization processes and time-averaging introduced during enamel sampling (e.g., van Dam and Reichart, 2009). The damping factor (d_f) is defined as the difference between the amplitude in the drinking water ($\delta^{18}\text{O}_{\text{mw}}$) and the measured amplitude in the tooth enamel ($\delta^{18}\text{O}_P$) relative to the amplitude in the drinking water (Passey and Cerling, 2002; Kohn, 2004). To correct for this signal loss in this study, we used a d_f of 0.5 for red deer and 0.6 for buffalo, following Kohn (2004).

Comparison of measured modern precipitation amounts and temperatures with those inferred from the stable isotope compositions of modern animal tooth enamel can provide a useful test of the paleoclimate relationships described above. Because no such data are available for our study area, we used stable isotope results




Fig. 6. (a) Monthly variations in air temperature (°C), rainfall $\delta^{18}O$ (‰, VSMOW) and rainfall amount (mm) in Zhengzhou city (~250 km from Luanchuan County). There is a negative shift in the $\delta^{18}O$ of precipitation during summer months (May–September) because of the rainfall amount effect. (b) Correlation of rainfall $\delta^{18}O$ with annual temperature, summer (April–September) and winter (October–March) temperatures. There is no significant correlation of rainfall $\delta^{18}O$ with summer half-year temperature due to the rainfall amount effect.

for modern buffalo from the Shunshanji archaeological site (33°34'58"N, 118°10'8"E) in Sihong County (Tian et al., 2013), which is located 700 km to the east. In that study, $\delta^{13}\text{C}_c$ and $\delta^{18}\text{O}_c$ were measured for ten subsamples taken from a premolar tooth at an interval of 5 mm along a growth band. The $\delta^{13}\text{C}_c$ values (-2.5 to -4.7‰) indicated that the buffalo had ingested a substantial amount of C_4 plants (Tian et al., 2013). As such, these results are unfortunately not suitable for rainfall reconstruction. The $\delta^{18}\text{O}_c$ values, however, showed typical seasonal variation (Fig. S2) and were used to estimate MAT, winter half-year and summer half-year temperatures. Before temperature reconstruction, $\delta^{18}\text{O}_c$ was translated to $\delta^{18}\text{O}_p$ using the average $\delta^{18}\text{O}_c - \delta^{18}\text{O}_p$ offset (7.7‰) for buffalo obtained in our study (Table 1). The reconstructed MAT using equation (4) is 13.0 °C (Table S2), which is comparable to the present mean annual temperature of 14.6 °C at that site. The inferred winter half-year temperature is 4.0 °C , which is $\sim 2\text{ °C}$ lower than measured directly. The inferred summer half-year temperature of 22.1 °C is very close to the measured value (22.9 °C) (Table S2). The general comparability between the inferred and measured temperatures provides a measure for the reliability of the temperature equations presented here. The negative offset of $\sim 2\text{ °C}$ for the inferred temperatures provides a useful reference when comparing inferred paleotemperatures with present temperatures at our study site.

The winter and summer $\delta^{18}\text{O}_p$, reconstructed $\delta^{18}\text{O}_{\text{mw}}$ and resulting average seasonal temperatures are listed in Table 3. The reconstructed winter (October–March) half-year temperatures varied from $-2.5 \pm 0.8\text{ °C}$ to $5.3 \pm 0.7\text{ °C}$ (mean: 0 ± 3 (1SD) °C), with the general condition being lower than the present temperature of $\sim 5\text{ °C}$ during the same months, notwithstanding the $\sim 2\text{ °C}$ offset observed for the winter temperature reconstruction using the modern buffalo tooth enamel. The inferred summer (April–September) temperatures showed a large range of variation from 8.4 ± 1.3 to $23.6 \pm 1.2\text{ °C}$ (mean: 15 ± 6 (1SD) °C). The inferred mean summer temperature is below the present average summer temperature of $\sim 20\text{ °C}$ for the same period in Luanchuan County.

If we examine the temperature data by sub-layer, a possible reason for the large variations becomes apparent. The inferred MAT, winter and summer temperatures for sub-layers 6 and 7 were

11.3 °C and 13.2 °C , 0 °C and 5.3 °C , and 23.6 °C and 21.3 °C , respectively. In contrast, the inferred MAT, winter and summer temperatures for sub-layer 5 were 3.4 °C to 7.9 °C , -2.5 °C to 0.7 °C , and 8.4 °C – 15.1 °C , respectively. The inferred temperatures for sub-layers 6 and 7 are much higher than those for sub-layer 5. Moreover, the inferred temperatures show a decreasing trend from sub-layer 7 to 5. Assuming a normal age–depth model, this change may represent a transition from interstadial to stadial mode, possibly Greenland Interstadial (GIS) 5 to its following stadial during ~ 33 – 31 cal ka BP as recorded in the $\delta^{18}\text{O}$ records of Chinese stalagmites and the GISP2 ice core (Wang et al., 2001). If so, GIS 5 had temperature conditions similar to today whereas the stadial was characterized by a mean annual temperature drop of $\sim 8\text{ °C}$. Our calculations also suggest that the climatic seasonality in Luanchuan County during the late MS 3 was variable, i.e., GIS 5 possessed a seasonality similar to present-day whereas the stadial had weaker seasonality than today.

5.4. Late MIS 3 paleoclimate estimates using different proxies in North China

Only a few previous studies have attempted to quantify paleoclimatic during the MIS 3 in North China. Li et al. (2011) reconstructed climatic change in Tangshan over the last 200 ka using a pollen-climate transfer function; the inferred MAT and MAP during 40–20 ka B.P. were lower by 5 °C and 150 mm , respectively, than present conditions. By comparison, Lu et al. (2007) estimated changes in MAT and MAP in Weinan in the southern part of the Chinese Loess Plateau over last 136 ka B.P. using phytolith assemblages. Their estimated values for MAT and MAP during 46–23 ka B.P. were 1 – 2 °C and 100 – 150 mm higher than those in the region today. Ning et al. (2008) reconstructed paleorainfall in Weinan and Lantian in Chinese Loess Plateau using the stable carbon isotope compositions of soil organic matter. They obtained estimates of MAP at 33 ka B.P. for both study sites that were $>200\text{ mm}$ lower than present conditions.

The difference in inferred MAP for Weinan obtained in the two previous studies can likely be attributed to the use of different proxy indicators, which have different sensitivities to temperature

Table 3
Calculated winter and summer average temperature based on mean tooth enamel $\delta^{18}\text{O}_p$ of red deer and buffalo.

Sample No.	Species	^a Winter $\delta^{18}\text{O}_p$ (‰, VSMOW)	^a Summer $\delta^{18}\text{O}_p$ (‰, VSMOW)	^b Winter $\delta^{18}\text{O}_p$ corrected (‰, VSMOW)	^b Summer $\delta^{18}\text{O}_p$ corrected (‰, VSMOW)	^c Winter $\delta^{18}\text{O}_{\text{mw}}$ (‰, VSMOW)	^c Summer $\delta^{18}\text{O}_{\text{mw}}$ (‰, VSMOW)	^d Winter T_{air} (Eq. (6)) (°C)	^e Summer T_{air} (from 2MAT–Winter T) (°C)
LCMD2(5):34	Red deer	+12.6 (n = 1)	+14.4 (n = 1)	+11.7 (n = 1)	+15.3 (n = 1)	–12.2	–9.1	0.7 ± 0.5	15.1 ± 1.1
LCMD2(6):19	Red deer	+13.9 (n = 1)	+19.3 (n = 1)	+11.2 (n = 1)	+22.0 (n = 1)	–12.7	–3.2	0 ± 0.5	23.6 ± 1.2
LCMA2(5):56-3	Buffalo	+12.2 ± 0.4 (n = 4)	+14.6 ± 0.1 (n = 2)	+10.4 ± 0.4 (n = 4)	+16.4 ± 0.1 (n = 2)	–13.4 ± 0.4	–8.1 ± 0.1	-1.0 ± 0.7	12.0 ± 1.2
LCMD2(5):16M1	Buffalo	–	–	–	–	–	–	–	–
LCMD2(5):16M2	Buffalo	+11.3 ± 0.5 (n = 3)	+14.1 (n = 1)	+9.2 ± 0.5 (n = 3)	+16.2 (n = 1)	–14.4 ± 0.5	–8.3	-2.5 ± 0.8	11.3 ± 1.3
LCMD2(5):16M3	Buffalo	+11.4 ± 0.7 (n = 3)	+13.4 ± 0.3 (n = 3)	+9.9 ± 0.7 (n = 3)	+14.9 ± 0.3 (n = 3)	–13.8 ± 0.7	–9.4 ± 0.3	-1.6 ± 1.0	8.4 ± 1.3
LCMD2(7):23-2	Buffalo	+16.0 ± 0.2 (n = 5)	+17.1 ± 0.1 (n = 9)	+15.2 ± 0.2 (n = 5)	+18.0 ± 0.1 (n = 9)	–9.1 ± 0.2	–6.7 ± 0.1	5.3 ± 0.7	21.3 ± 1.4

^a The mean winter and summer $\delta^{18}\text{O}_p$ were respectively calculated based on the lowest and highest values of seasonal $\delta^{18}\text{O}_p$ curve from each tooth (as indicated in Fig. 3). The \pm on each average stands for actual range.

^b Winter $\delta^{18}\text{O}_p$ corrected = (Summer $\delta^{18}\text{O}_p$ + Winter $\delta^{18}\text{O}_p$)/2 - (Summer $\delta^{18}\text{O}_p$ - Winter $\delta^{18}\text{O}_p$)/2d_r; Summer $\delta^{18}\text{O}_p$ corrected = (Summer $\delta^{18}\text{O}_p$ + Winter $\delta^{18}\text{O}_p$)/2 + (Summer $\delta^{18}\text{O}_p$ - Winter $\delta^{18}\text{O}_p$)/2d_r. Values of d_r are 0.5 for red deer and 0.6 for buffalo.

^c Winter and summer $\delta^{18}\text{O}_{\text{mw}}$ were calculated based on winter $\delta^{18}\text{O}_p$ corrected and summer $\delta^{18}\text{O}_p$ corrected, respectively, using equation (2) for buffalo and equation (3) for red deer.

^d Winter T_{air} was calculated winter $\delta^{18}\text{O}_{\text{mw}}$ using equation (6). The \pm represents an overall uncertainty that takes the variation in winter $\delta^{18}\text{O}_{\text{mw}}$ and the uncertainty of equation (6) into consideration.

^e Summer T_{air} was calculated by subtracting the winter T_{air} from 2MAT (eq. (4)). The \pm was obtained according to error propagation equation for the calculation.

and rainfall. The pollen and phytolith assemblages both provide an indication of changes in plant compositions, which tend to represent long-term changes in regional climate. These two proxy indicators also reflect overall environmental conditions and cannot be used to differentiate between the effects of temperature and rainfall. By comparison, paleorainfall reconstructions using the carbon isotope of soil organic matter should be relatively reliable because the $\delta^{13}\text{C}$ of terrestrial higher plants is much more responsive to changes in rainfall than temperature (Liu et al., 2005). The work of Ning et al. (2008), therefore, which suggests a dry late MIS 3, is likely more reliable than estimates from the pollen and phytolith assemblages. That said, the inferred rainfall amounts using soil organic matter $\delta^{13}\text{C}$ provides only a long-term measure of local conditions due to the limited time resolution of samples from loess-paleosol sections.

The $\delta^{13}\text{C}_c$ and $\delta^{18}\text{O}_p$ of tooth enamel are independent proxies that can be used to reliably infer rainfall and temperature, respectively, on an inter- or intra-annual basis, depending on sampling resolution. In contrast to the warm and wet conditions during MIS 3 proposed in previous studies (e.g., Li et al., 2003; Lu et al., 2007; Chen and Wu, 2008; Yang et al., 2011; Li et al., 2014), our results indicate a generally colder and drier late MIS 3 at our study location. Such dry conditions are consistent with long-term records inferred from soil organic carbon isotope records for the Chinese Loess Plateau during late MIS 3 (Ning et al., 2008).

Solar insolation intensity and global ice volume are two main forcing factors of Earth's climate. During the late MIS 3, average solar insolation over the summer months (June–August) at 33°N , where the Longquan Cave lies, was slightly higher ($\sim 10.1 \text{ W/m}^2$) than today (Berger, 1978; Laskar et al., 2004). This condition should facilitate relatively warm overall conditions in the Northern Hemisphere. That said, millennial-scale abrupt climatic events (i.e., Dansgaard-Oeschger cycles and Heinrich events) were common during the MIS 3 and have been well documented in stalagmite $\delta^{18}\text{O}$ records for Central China (Wang et al., 2001). In North China, abrupt climatic events such as Heinrich 1 and 2, characterized by cold and dry conditions, have also been recognized in studies of loess deposits (e.g., Wang et al., 2012).

As aforementioned, the decrease in the inferred temperature from sub-layer 7 to 5 may represent a transition from GIS 5 to its following stadial at $\sim 33\text{--}31 \text{ ka}$. This change may explain our inferred lower MAT and MAP at Luanchuan County at this time. We cannot be certain whether the tooth enamel isotopic results capture the full range of climatic change during this period but they at least show $\sim 8^\circ\text{C}$ variation on an annual time-scale over the interstadial-stadial cycle. Even within sub-layer 5, we observe large inter-annual variations both in MAT and seasonal temperature differences, which may indicate climatic instability on an annual time-scale during the stadial.

An expanded global ice volume may also have contributed to the relatively dry late MIS 3 climate in north China. Ice sheets in the Northern Hemisphere gradually expanded from MIS 3 to the Last Glacial Maximum (LGM) at a rate of $1\text{--}2 \text{ cm/y}$ (Thompson and Goldstein, 2005), causing a continuous drop in sea level. Sea level during the late MIS 3 was $\sim 60\text{--}90 \text{ m}$ lower than today (Chappell, 2002; Siddall et al., 2003). Expansion of the ice sheets in Northern Hemisphere during the last MIS 3 could have strengthened polar high pressure, which would have weakened the thermal contrast between continents and the ocean, eventually leading to a decrease in rainfall amount (Yang and Ding, 2008).

6. Conclusions

Stable carbon and oxygen isotope compositions of fossil ($\sim 33\text{--}31 \text{ ka B.P.}$) mammal tooth enamel (buffalo and red deer)

recovered from Longquan Cave in North China were analyzed to decipher the paleoecology and paleoenvironment at this site during the late MIS 3. These data support the following conclusions:

- (1) The $\delta^{13}\text{C}$ of plants in the diet of these animals, as calculated from bioapatite structural carbonate, ranged from -29 to -24‰ , indicating a diet of C_3 plants for both species.
- (2) The amount of mean annual precipitation was estimated using the relationship between modern $\delta^{13}\text{C}$ of C_3 vegetation and MAP in North China. The estimated annual precipitation during MIS 3 at our study site ranged from 363 to 670 mm (mean: $593 \pm 104 \text{ mm}$), which is substantially lower than the annual average for modern precipitation (873 mm) in the region.
- (3) The oxygen isotope compositions of bioapatite phosphate ($\delta^{18}\text{O}_p$) along growth bands of tooth enamel were used to reconstruct mean annual and seasonal air temperatures, based on the present-day relationships between $\delta^{18}\text{O}_p$ and meteoric water ($\delta^{18}\text{O}_{\text{mw}}$) and between $\delta^{18}\text{O}_{\text{mw}}$ and air temperature. The reconstructed MAT ranged from 3 to 13°C (mean of $8 \pm 4.0^\circ\text{C}$), which is lower than the present average annual temperature (12°C). Calculated summer (April–September) and winter (October–March) temperatures were $8\text{--}24^\circ\text{C}$ (mean: $15 \pm 6^\circ\text{C}$) and -3 to 5°C (mean: $0 \pm 3^\circ\text{C}$), respectively, suggesting either similar or relatively weaker seasonality at the study site during the late MIS 3 compared to the present summer-winter contrast of $\sim 15^\circ\text{C}$.

Based on the above results, the climate at Longquan Cave, North China, during the late MIS 3 was characterized by relatively cold and dry conditions. This observation contrasts with long-term warm and wet conditions suggested in some previous studies. Our data, however, point to generally cold and dry but also fluctuating climatic conditions on an annual timescale. This result may signal an abrupt climatic cooling event from Greenland Interstadial Stage (GIS) 5 to its following stadial. The climate at GIS 5 may have been as warm as today whereas the following stadial was $\sim 8^\circ\text{C}$ colder than GIS 5.

Author contribution statement

Xu Wang: Project administration, Funding acquisition, Conceptualization, Writing-Original Draft. Jing Sun: Data curation, Formal analysis, Visualization. Fred J. Longstaffe: Data curation, Formal analysis, Funding acquisition, Writing-Review & Editing. Xuejun Gu: Resources. Shuisheng Du: Resources, Writing-Review & Editing. Linlin Cui: Data curation, Formal analysis. Xiaozhong Huang: Data curation, Formal analysis. Zhongli Ding: Writing-Review & Editing.

Declaration of competing interest

We declare that we have no competing interests.

Acknowledgements

This research was financially supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDB26000000 and XDA19050104), the Key Research Program of the Institute of Geology and Geophysics (Grant No. IGGCAS-201905). Many of the phosphate oxygen isotope analyses were financially supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant program (FJL) (Grant No.05904-2019 RGPIN). This is the University of Western Ontario's Laboratory for Stable Isotope Science (LSIS) Contribution #383.

Grace Yau (LSIS) is thanked for her technical assistance.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.quascirev.2021.107222>.

References

- Amiot, R., Lécuyer, C., Buffetaut, E., Fluteau, F., Legendre, S., Martineau, F., 2004. Latitudinal temperature gradient during the cretaceous upper campanian middle maastrichtian: $\delta^{18}\text{O}$ record of continental vertebrates. *Earth Planet Sci. Lett.* 226, 255–272.
- Amiot, R., Wang, X., Zhou, Z., Wang, X.L., Buffetaut, E., Lécuyer, C., Ding, Z., Fluteau, F., Kellner, A.W.A., Tong, H.Y., Zhang, F., 2010. Oxygen and carbon isotope compositions of middle Cretaceous vertebrates from North Africa and Brazil: ecological and environmental significance. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 297, 439–451.
- Amiot, R., Wang, X., Zhou, Z., Wang, X.L., Buffetaut, E., Lécuyer, C., Ding, Z., Fluteau, F., Hibino, T., Kusuhashi, N., Mo, J.Y., Suteethorn, V., Wang, Y.Q., Xu, X., Zhang, F.S., 2011. Oxygen isotopes of East Asian dinosaurs reveal exceptionally cold early cretaceous climates. *Proceedings of National Academy of Sciences of USA* 108, 5179–5183.
- Amiot, R., Wang, X., Zhou, Z., Wang, X.L., Lécuyer, C., Buffetaut, E., Fluteau, F., Ding, Z., Kusuhashi, N., Mo, J.Y., Philippe, M., Suteethorn, V., Wang, Y.Q., Xu, X., 2015. Environment and ecology of East Asian dinosaurs during the Early Cretaceous inferred from stable oxygen and carbon isotopes in apatite. *J. Asian Earth Sci.* 98, 358–370.
- Azorit, C., Analla, M., Carrasco, R., Calvo, J.A., Muñoz-Cobo, J., 2002. Teeth eruption pattern in red deer (*Cervus elaphus hispanicus*) in southern Spain. *Ann. Biol.* 24, 107–114.
- Bendrey, R., Vella, D., Zazzo, A., Balasse, M., Lepetz, S., 2015. Exponentially decreasing tooth growth rate in horse teeth: implications for isotopic analyses. *Archaeometry* 57, 1104–1124.
- Berger, A., 1978. Long-term variations of daily insolation and Quaternary climate changes. *J. Atmos. Sci.* 35, 2362–2367.
- Bernard, A., Daux, V., Lécuyer, C., Brugal, J.P., Genty, D., Wainer, K., Gardien, V., Fourel, F., Jaubert, J., 2009. Pleistocene seasonal temperature variations recorded in the $\delta^{18}\text{O}$ of *Bison priscus* teeth. *Earth Planet Sci. Lett.* 283, 133–143.
- Bonafini, M., Pellegrini, M., Ditchfield, P., Pollard, A.M., 2013. Investigation of the 'canopy effect' in the isotope ecology of temperate woodlands. *J. Archaeol. Sci.* 40, 3926–3935.
- Brown, W.A.B., Chapman, N.G., 1991. Age assessment of red deer (*Cervus elaphus*): from a scoring scheme based on radiographs of developing permanent molariform teeth. *J. Zool.* 225, 85–97.
- Bryant, J.D., Luz, B., Froelich, P.N., 1994. Oxygen isotopic composition of fossil horse tooth phosphate as a record of continental paleoclimate. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 107, 303–316.
- Bryant, D., Froelich, P., Showers, W., Genna, B., 1996. Biologic and climatic signals in the oxygen isotopic composition of Eocene–Oligocene equid enamel phosphate. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 126, 75–89.
- Cerling, T.E., Harris, J., MacFadden, B., Leakey, M., Quade, J., Eisenmann, V., Ehleringer, J., 1997. Global vegetation change through the Miocene–Pliocene boundary. *Nature* 389, 153–158.
- Cerling, T.E., Harris, J.M., 1999. Carbon isotope fractionation between diet and bioapatite in ungulate mammals and implications for ecological and paleoecological studies. *Oecologia* 120, 347–363.
- Chappell, J., 2002. Sea level changes forced ice breakouts in the Last Glacial Cycle: new results from coral terraces. *Quat. Sci. Rev.* 21, 1229–1240.
- Chen, X.Y., Wu, N.Q., 2008. Relatively warm-humid climate recorded by mollusk species in the Chinese Loess Plateau during MIS 3 and its possible forcing mechanism. *Quat. Sci.* 28, 154–161 (In Chinese with English abstract).
- Compilatory Commission of Annals of Luanchuan County, 1994. In: *Annals of Luanchuan County*. SDX Joint Publishing Company, Beijing, pp. 74–77.
- Coplen, T.B., Kendall, C., Hoppie, J., 1983. Comparison of stable isotope reference samples. *Nature* 302, 236–238.
- Cormie, A.B., Luz, B., Schwarcz, H.P., 1994. Relationship between the hydrogen and oxygen isotopes of deer bone and their use in the estimation of relative humidity. *Geochem. Cosmochim. Acta* 58, 3439–3449.
- Crowson, R.A., Showers, W.J., Wright, E.K., Hoering, T.C., 1991. A method for preparation of phosphate samples for oxygen isotope analysis. *Anal. Chem.* 63, 2397–2400.
- D'Angela, D., Longinelli, A., 1990. Oxygen isotopes in living mammal's bone phosphate: further results. *Chem. Geol.* 86, 75–82.
- Dansgaard, W., 1964. Stable isotopes in precipitation. *Tellus* 16, 436–468.
- Dansgaard, W., Johnsen, S., Clausen, H.B., Dahl-Jensen, D., Gundestrup, N., Hammer, C.U., Oeschger, H., 1984. North Atlantic climatic oscillations revealed by deep Greenland ice cores. In: Hansen, J.E., Takahashi, T. (Eds.), *Climate Processes and Climate Sensitivity*. American Geophysical Union, Washington, pp. 288–298.
- Dansgaard, W., Johnsen, S.J., Clausen, H.B., Dahl-Jensen, D., Gundestrup, N.S., Hammer, C.U., Hvidberg, C.S., Steffensen, J.P., Sveinbjörnsdóttir, A.E., Jouzel, J., Bond, G., 1993. Evidence for general instability of past climate from a 250-kyr ice-core record. *Nature* 364, 218–220.
- Deines, P., 1980. The isotopic composition of reduced organic carbon. In: Fritz, P., Fontes, J.C. (Eds.), *Handbook of Environmental Isotope Geochemistry*. Elsevier, Amsterdam, pp. 329–406.
- Delgado Huertas, A., Iacumin, P., Longinelli, A., 1997. A stable isotope study of fossil mammal remains from the Paglicci cave, southern Italy, 13 to 33 ka BP: palaeoclimatological considerations. *Chem. Geol.* 141, 211–223.
- Deng, T., Dong, J., Wang, Y., 2002. Variation of terrestrial ecosystem recorded by stable carbon isotopes of fossils in northern China during the Quaternary. *Chin. Sci. Bull.* 47 (1), 76–78.
- Deng, T., Li, Y.M., 2005. Vegetational ecotype of the Gyirong Basin in Tibet, China and its response in stable carbon isotopes of mammal tooth enamel. *Chin. Sci. Bull.* 50 (12), 1225–1229.
- Dettman, D.L., Kohn, M.J., Quade, J., Ryerson, F.J., Ojha, T.P., Hamidullah, S., 2001. Seasonal stable isotope evidence for a strong Asian monsoon throughout the past 10.7 m.y. *Geology* 29, 31–34.
- Drucker, D.G., Bridault, A., Hobson, K.A., Szuma, E., Bocherens, H., 2008. Can carbon-13 in large herbivores reflect the canopy effect in temperate and boreal ecosystems? Evidence from modern and ancient ungulates. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 266, 69–82.
- Du, S.S., Li, X., Zhou, L., Pang, H.J., Bar-Yosef, O., Wu, X.H., 2016. Longquan cave: an early upper palaeolithic site in Henan province, China. *Antiquity* 90, 876–893.
- Fabre, M., Lécuyer, C., Brugal, J.-P., Amiot, R., Fourel, F., Martineau, F., 2011. Late Pleistocene climatic change in the French Jura (Gigny) recorded in the $\delta^{18}\text{O}$ of phosphate from ungulate tooth enamel. *Quat. Res.* 75, 605–613.
- Faegri, K., Kaland, P.E., Krzywinski, K., 1989. In: *Textbook of Pollen Analysis*, fourth ed. John Wiley & Sons, London, pp. 1–328.
- Farquhar, G., Ehleringer, K., Hubick, K., 1989. Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40, 503–537.
- Flanagan, L.B., Farquhar, G.D., 2014. Variation in the carbon and oxygen isotope composition of biomass and its relationship to water-use efficiency at the leaf- and ecosystem-scales in a northern Great Plains grassland. *Plant Cell Environ.* 37, 425–438.
- Francey, R.J., Farquhar, G.D., 1982. An explanation of $^{13}\text{C}/^{12}\text{C}$ variations in tree rings. *Nature* 295, 28–31.
- Fricke, H.C., O'Neil, J.R., 1996. Inter- and intra-tooth variation in the oxygen isotope composition of mammalian tooth enamel phosphate: implications for palaeoclimatological and palaeobiological research. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 126, 91–99.
- Fricke, H.C., O'Neil, J.R., 1999. The correlation between $^{18}\text{O}/^{16}\text{O}$ ratio of meteoric water and surface temperature: its use in investigating terrestrial climate change over geologic time. *Earth Planet Sci. Lett.* 170, 181–196.
- Gadbury, C., Todd, L., Jahren, A.H., Amundson, R., 2000. Spatial and temporal variations in the isotopic composition of bison tooth enamel from the Early Holocene Hudson-Meng Bone Bed, Nebraska. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 157, 79–93.
- Gibb, O.T., Hillaire-Marcel, C., de Vernal, A., 2014. Oceanographic regimes in the northwest Labrador Sea since Marine Isotope Stage 3 based on dinocyst and stable isotope proxy records. *Quat. Sci. Rev.* 92, 269–279.
- Godawa, J., 1989. Age determination in the Red deer (*Cervus elaphus*). *Acta Theriol.* 34, 381–384.
- Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the Northeast Atlantic Ocean during the past 130,000 years. *Quat. Res.* 29, 142–152.
- Hoppe, K.A., 2006. Correlation between the oxygen isotope ratio of North American bison teeth and local waters, implication for paleoclimatic reconstructions. *Earth Planet Sci. Lett.* 244, 408–417.
- Iacumin, P., Bocherens, H., Mariotti, A., Longinelli, A., 1996. Oxygen isotope analyses of co-existing carbonate and phosphate in biogenic apatite: a way to monitor diagenetic alteration of bone phosphate? *Earth Planet Sci. Lett.* 142, 1–6.
- Imbrie, J., Hays, J.D., Martinson, D.G., McIntyre, A., Mix, A.C., Morley, J.J., Pisias, N.G., Prell, W.L., Shackleton, N.J., 1984. The orbital theory of Pleistocene climate: support from a revised chronology of the marine $\delta^{18}\text{O}$ record. In: Berger, A., Imbrie, J., Hays, J., Kukla, G., Saltzman, B. (Eds.), *Milankovitch and Climate (Part 1)*. Dordrecht Reidel Publishing, Dordrecht, pp. 269–305.
- Jiang, H.C., Ding, Z.L., 2005. Temporal and spatial changes of vegetation cover on the Chinese Loess Plateau through the last glacial cycle: evidence from spore-pollen records. *Rev. Palaeobot. Palynol.* 133, 23–37.
- Jiménez-Moreno, G., Anderson, R.S., Desprat, S., Grigg, L.D., Grimm, E.C., Heusser, L.E., Jacobs, B.F., López-Martínez, C., Whitlock, C.L., Willard, D.A., 2010. Millennial-scale variability during the last glacial in vegetation records from North America. *Quat. Sci. Rev.* 29, 2865–2881.
- Johnsen, S.J., Clausen, H.B., Dansgaard, W., Fuhrer, K., Gundestrup, N., Hammer, C.U., Iversen, P., Jouzel, J., Stauffer, B., Steffensen, J.P., 1992. Irregular glacial interstadials recorded in a new Greenland ice core. *Nature* 359, 311–313.
- Keeling, C.D., 1979. The Suess effect: ^{13}C – ^{14}C interrelations. *Environ. Int.* 2 (4/6), 229–300.
- Koch, P.L., Tuross, N., Fogel, M.L., 1997. The effects of sample treatment and diagenesis on the isotopic integrity of carbonate in biogenic hydroxylapatite. *J. Archaeol. Sci.* 24, 417–429.
- Koch, P., 1998. Isotopic reconstruction of past continental environments. *Annu. Rev. Earth Planet Sci.* 26, 573–613.
- Koch, P.L., Hoppe, K.A., Webb, S.D., 1998. The isotopic ecology of late Pleistocene mammals in North America. Part 1. Florida. *Chem. Geol.* 152, 119–138.
- Kohn, M.J., Schoeninger, M.J., Valley, J.W., 1996. Herbivore tooth oxygen isotope

- compositions: effects of diet and physiology. *Geochem. Cosmochim. Acta* 60, 3889–3896.
- Kohn, M.J., Cerling, T.E., 2002. Stable isotope compositions of biological apatite. In: Kohn, M., Rakovan, J., Hughes, J. (Eds.), *Phosphates—Geochemical, Geobiological, and Materials Importance. Reviews in Mineralogy and Geochemistry*, vol. 48. Mineralogical Society of America, Washington D.C., pp. 455–488.
- Kohn, M.J., 2004. Comment: tooth enamel mineralization in ungulates: implications for recovering a primary isotopic time-series, by B.H. Passey and T.E. Cerling (2002). *Geochem. Cosmochim. Acta* 68, 403–405.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M., Levrard, B., 2004. A long-term numerical solution for the insolation quantities of the Earth. *Astron. Astrophys.* 428, 261–285.
- Lécuyer, C., Grandjean, P., O'Neil, J.R., Cappetta, H., Martineau, F., 1993. Thermal excursions in the ocean at the Cretaceous-Tertiary boundary (northern Morocco): $\delta^{18}\text{O}$ record of phosphatic fish debris. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 105, 235–243.
- Li, Y.M., Liu, T.S., Wu, W.X., Han, J.M., Hong, Y.T., 2003. Palaeoenvironment in Chinese Loess Plateau during MIS 3: evidence from malan loess. *Quat. Sci.* 23, 69–76 (In Chinese with English abstract).
- Li, Y.S., Chen, J.Q., Zhao, S., Wang, H.Y., Li, X., 2011. Quantitative reconstruction of palaeoclimate since 200 ka BP in Tangshan area, Hebei Province. *Mar. Geol. Quat. Geol.* 31, 163–170 (In Chinese with English abstract).
- Li, X.Q., Gao, Q., Hou, Y.M., Zhao, K.L., Sun, N., Yang, Z.M., Zhen, Z.M., Liu, Y., Zhou, X.Y., 2014. The vegetation and environment at the Wulumulun site in the Ordos Plateau, Inner Mongolia during MIS 3 period. *Acta Anthropol. Sin.* 33, 60–69.
- Liu, W.G., Feng, X.H., Ning, Y.F., Zhang, Q.L., Cao, Y.N., 2005. $\delta^{13}\text{C}$ variation of C_3 and C_4 plants across an Asian monsoon rainfall gradient in arid northwestern China. *Global Change Biol.* 11, 1094–1100.
- Longinelli, A., 1984. Oxygen isotopes in mammal bone phosphate: a new tool for paleohydrological and paleoclimatological research? *Geochem. Cosmochim. Acta* 48, 385–390.
- Lourantou, A., Lavric, J.V., Köhler, P., Barnola, J.-M., Paillard, D., Michel, E., Raynaud, D., Chappellaz, J., 2010. Constraint of the CO_2 rise by new atmospheric carbon isotopic measurements during the last deglaciation. *Global Biogeochem. Cycles* 24, GB2015. <https://doi.org/10.1029/2009GB003545>.
- Lu, H.Y., Wu, N.Q., Liu, K.B., Jiang, H., Liu, T.S., 2007. Phytoliths as quantitative indicators for the reconstruction of past environmental conditions in China II: palaeoenvironmental reconstruction in the Loess Plateau. *Quat. Sci. Rev.* 26, 759–772.
- Luz, B., Kolodny, Y., Horowitz, M., 1984. Fractionation of oxygen isotopes between mammalian bone-phosphate and environmental drinking water. *Geochem. Cosmochim. Acta* 48, 1689–1693.
- MacFadden, B., Cerling, T., Prado, J., 1996. Cenozoic terrestrial ecosystem evolution in Argentina: evidence from carbon isotopes of fossil mammal teeth. *Palaios* 11, 319–327.
- Maher, L.J., 1981. Statistics for microfossil concentration measurements employing samples spiked with marker grains. *Rev. Palaeobot. Palynol.* 32, 153–191.
- Marino, B.D., McElroy, M.B., Salawitch, R.J., Spaulding, W.G., 1992. Glacial-to-interglacial variations in the carbon isotopic composition of atmospheric CO_2 . *Nature* 357, 461–466.
- Metcalfe, J.Z., Longstaffe, F.J., Ballenger, J.A.M., Vance Haynes Jr., C., 2011. Isotopic paleoecology of Clovis mammoths from Arizona. *Proceedings of National Academy of Sciences of USA* 108 (44), 17916–17920.
- Moss-Salentijn, L., Moss, M.L., Yuan, M.S., 1997. The ontogeny of mammalian enamel. In: Koeningswald, W.V., Sander, P.M. (Eds.), *Tooth Enamel Microstructure*. A. A. Balkema, pp. 5–30.
- Ning, Y.F., Liu, W.G., An, Z.S., 2008. A 130-ka reconstruction of precipitation on the Chinese Loess Plateau from organic carbon isotopes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 270, 59–63.
- O'Leary, M.H., 1981. Carbon isotope fractionation in plants. *Phytochemistry* 20, 553–567.
- O'Leary, M.H., 1988. Carbon isotopes in photosynthesis. *Bioscience* 38, 328–336.
- Passey, B.H., Cerling, T.E., 2002. Tooth enamel mineralization in ungulates: implications for recovering a primary isotopic time-series. *Geochem. Cosmochim. Acta* 66, 3225–3234.
- Passey, B.H., Robinson, T.F., Ayliffe, L.K., Cerling, T.E., Sponheimer, M., Dearing, M.D., Roeder, B.L., Ehleringer, J.R., 2005. Carbon isotope fractionation between diet, breath CO_2 , and bioapatite in different mammals. *J. Archaeol. Sci.* 32, 1459–1470.
- Pellegrini, M., Snoeck, C., 2016. Comparing bioapatite carbonate pre-treatments for isotopic measurements: Part 2- Impact on carbon and oxygen isotope compositions. *Chem. Geol.* 420, 88–96.
- Quade, J., Cerling, T.E., Barry, J.C., Morgan, M.M., Pilbeam, D.R., Chivas, A.R., Lee-Thorp, J.A., van der Merwe, N.J., 1992. A 16 million year record of paleodiet from Pakistan using carbon isotopes in fossil teeth. *Chem. Geol.* 94, 183–192.
- Rao, Z.G., Jia, G.D., Zhu, Z.Y., Wu, Y., Zhang, J.W., 2008. Comparison of carbon isotope composition of total organic carbon and long-chain n-alkanes from surface soils in eastern China and their significance. *Chin. Sci. Bull.* 53, 3921–3927.
- Rozanski, K., Araguas-Araguas, L., Gonfiantini, R., 1993. Isotopic patterns in modern global precipitation. In: Swart, P., Lohmann, K., McKenzie, J., Savin, S. (Eds.), *Climate Change in Continental Isotopic Records. Geophysical Monograph*, vol. 78. American Geophysical Union, Washington, DC, pp. 1–36.
- Sakai, T., Hirai, G., 1982. Calcification and crystallization in bovine enamel. *J. Dent. Res.* 61, 57–59.
- Sánchez Goñi, M.F., Cacho, I., Turon, J.-L., Guiot, J., Sierro, F.J., Peyrouquet, J.-P., Grimalt, J.O., Shackleton, N.J., 2002. Synchronicity between marine and terrestrial responses to millennial scale climatic variability during the last glacial period in the Mediterranean region. *Clim. Dynam.* 19, 95–105.
- Sánchez Goñi, M.F., Harrison, S.P., 2010. Millennial-scale climate variability and vegetation changes during the last glacial: concepts and terminology. *Quat. Sci. Rev.* 29, 2823–2827.
- Schleser, G., Jayasekera, R., 1985. $\delta^{13}\text{C}$ variations of leaves in forests as an indication of reassimilated CO_2 from the soil. *Oecologia* 65, 536–542.
- Siddall, M., Rohling, E.J., Almogi-Labin, A., Hemleben, Ch., Meischner, D., Schmelzer, I., Smeed, D.A., 2003. Sea-level fluctuations during the last glacial cycle. *Nature* 423, 853–858.
- Sionneau, T., Bout-Roumazielles, V., Meunier, G., Kissel, C., Flower, B.P., Bory, A., Tribouillard, N., 2013. Atmospheric re-organization during marine isotope stage 3 over the North American continent: sedimentological and mineralogical evidence from the gulf of Mexico. *Quat. Sci. Rev.* 81, 62–73.
- Snoeck, C., Pellegrini, M., 2015. Comparing bioapatite carbonate pre-treatments for isotopic measurements: Part 1- Impact on structure and chemical composition. *Chem. Geol.* 417, 394–403.
- Spötl, C., Vennemann, T.W., 2003. Continuous-flow isotope ratio mass spectrometric analysis of carbonate minerals. *Rapid Commun. Mass Spectrom.* 17, 1004–1006.
- Sternberg, L., Mulkey, S., Wright, S., 1989. Ecological interpretation of leaf carbon isotope ratios: influence of respired carbon dioxide. *Ecology* 70, 1317–1324.
- Stevens, R.E., Balasse, M., O'Connell, T.C., 2011. Intra-tooth oxygen isotope variation in a known population of red deer: implications for past climate and seasonality reconstructions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 301, 64–74.
- Straight, W.H., Barrick, R.E., Eberth, D.A., 2004. Reflections of surface water, seasonality and climate in stable oxygen isotopes from tyrannosaurid tooth enamel. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 206, 239–256.
- Stuiver, M., Grootes, P.M., 2000. GISP2 oxygen isotope ratios. *Quat. Res.* 53, 277–284.
- Suga, S., 1979. Comparative histology of progressive mineralization pattern of developing incisor enamel of rodents. *J. Dent. Res.* 58, 1025–1026.
- Suga, S., 1982. Progressive mineralization pattern of developing enamel during the maturation stage. *J. Dent. Res.* 61, 1532–1542.
- Suga, S., Murayama, Y., Musashi, T., 1970. A study of the mineralization process in the developing enamel of Guinea pigs. *Arch. Oral Biol.* 15, 597–612.
- Suga, S., Aoki, H., Yamashita, Y., Tsuno, M., Ogawa, M., 1987. A comparative study of disturbed mineralization of rat incisor enamel induced by strontium and fluoride administration. *Adv. Dent. Res.* 1, 339–355.
- Thompson, W.G., Goldstein, S.L., 2005. Open-system coral ages reveal persistent suborbital sea-level cycles. *Science* 308, 401–404.
- Tian, X.S., Zhu, C., Shui, T., Huang, Y.P., 2013. Diets, eco-environments and seasonal variations recorded in the oxygen and carbon isotopic compositions of mammal tooth enamel from the Shunshanji site, Sihong county, Jiangsu Province, China. *Chin. Sci. Bull.* 58 (31), 3788–3795.
- Traylor, R.B., Kohn, M.J., 2016. Tooth enamel maturation reequilibrates oxygen isotope compositions and supports simple sampling methods. *Geochem. Cosmochim. Acta* 198, 32–47.
- Tütken, T., Furrer, H., Walter Vennemann, T., 2007. Stable isotope compositions of mammoth teeth from Niederweningen, Switzerland: implications for the Late Pleistocene climate, environment, and diet. *Quat. Int.* 164/165, 139–150.
- van Dam, J.A., Reichert, G.J., 2009. Oxygen and carbon isotope signatures in late Neogene horse teeth from Spain and application as temperature and seasonality proxies. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 274, 64–81.
- Van der Merwe, N., Medina, E., 1989. Photosynthesis and $^{13}\text{C}/^{12}\text{C}$ ratios in Amazon rain forests. *Geochem. Cosmochim. Acta* 53, 1091–1094.
- Van Meerbeek, C.J., Renssen, H., Roche, D.M., Wohlfarth, B., Bohncke, S.J.P., Bos, J.A.A., Engels, S., Helmens, K.F., Sánchez Goñi, M.F., Svensson, A., Vandenberghe, J., 2011. The nature of MIS 3 stadial-interstadial transitions in Europe: new insights from model-data comparisons. *Quat. Sci. Rev.* 30, 3618–3637.
- Wang, F.X., Qian, N.F., Zhang, Y.L., Yang, H.Q., 1995. In: *Pollen Flora of China*. Science Press, Beijing, p. 461 (In Chinese).
- Wang, G.A., Han, J.M., Liu, T.S., 2003. The carbon isotopic composition of C_3 herbaceous plants in loess area of North China. *Science in China (Series D)* 46, 1069–1076.
- Wang, G.A., Li, J.Z., Liu, X.Z., Li, X.Y., 2013b. Variations in carbon isotope ratios of plants across a temperature gradient along the 400 mm isohet of mean annual precipitation in north China and their relevance to paleovegetation reconstruction. *Quat. Sci. Rev.* 63, 83–90.
- Wang, X., Ding, Z.L., Peng, P.A., 2012. Changes in fire regimes in the Chinese Loess Plateau since the Last Glacial Maximum and implications for linkages to paleoclimate and past human activity. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 315/316, 61–74.
- Wang, X., Cui, L.L., Xiao, J.L., Ding, Z.L., 2013a. Stable carbon isotope of black carbon in lake sediments as an indicator of terrestrial environmental changes: an evaluation on paleorecord from Daihai Lake, Inner Mongolia, China. *Chem. Geol.* 347, 123–134.
- Wang, Y., Cerling, T.E., MacFadden, B., 1994. Fossil horses and carbon isotopes: new evidence for Cenozoic dietary, habitat, and ecosystem changes in North America. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 107, 269–279.
- Wang, Y., Deng, T., 2005. A 25 m.y. isotopic record of paleodiet and environmental change from fossil mammals and paleosols from the NE margin of the Tibetan

- Plateau. *Earth Planet Sci. Lett.* 236, 322–338.
- Wang, Y.J., Lv, H.Y., Wang, G.A., Yang, H., Li, Z., 2000. Carbon isotope analysis of silicates in C3, C4 plants and modern soils. *Chin. Sci. Bull.* 45, 978–982 (In Chinese).
- Wang, Y.J., Cheng, H., Edwards, R.L., An, Z.S., Wu, J.Y., Shen, C.-C., Dorale, J.A., 2001. A high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China. *Science* 294, 2345–2348.
- Webb, E.C., White, C.D., Longstaffe, F.J., 2014. Investigating inherent differences in isotopic composition between human bone and enamel bioapatite: implications for reconstructing residential histories. *J. Archaeol. Sci.* 50, 97–107.
- Wu, H., Hou, W., Qian, Z.H., Hu, J.G., 2012. The research on the sensitivity of climate change in China in recent 50 years based on composite index. *Acta Phys. Sin.* 61 (14), 149205.
- World Data Center for Paleoclimatology, 2006. Boulder and NOAA Paleoclimatology program [DB/OL]. http://hurricane.ncdc.noaa.gov/pls/paleox/f?p=519:1:::::P1_STUDY_ID:9972.
- Yang, Q., Li, X.Q., Zhou, X.Y., Liu, H.H., Zhao, K.L., Sun, N., 2011. Vegetation succession and its response to climate changes since MIS 3 in desert-loess transition belt, Northern China. *Quat. Sci.* 31 (6), 962–971 (In Chinese with English abstract).
- Yang, S.L., Ding, Z.L., 2008. Advance-retreat history of the East-Asian summer monsoon rainfall belt over northern China during the last two glacial–interglacial cycles. *Earth Planet Sci. Lett.* 274, 499–510.
- Yang, S.L., Ding, Z.L., Wang, X., Tang, Z.H., Gu, Z.Y., 2012. Negative $\delta^{18}\text{O}$ – $\delta^{13}\text{C}$ relationship of pedogenic carbonate from northern China indicates a strong response of C3/C4 biomass to the seasonality of Asian monsoon precipitation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 317/318, 32–40.
- Zazzo, A., Balasse, M., Patterson, W.P., 2005. High-resolution $\delta^{13}\text{C}$ intratooth profiles in bovine enamel: implications for mineralization pattern and isotopic attenuation. *Geochem. Cosmochim. Acta* 69, 3631–3642.
- Zazzo, A., Balasse, M., Passey, B.H., Moloney, A.P., Monahan, F.J., Schmidt, O., 2010. The isotope record of short- and long-term dietary changes in sheep tooth enamel: implications for quantitative reconstruction of paleodiets. *Geochem. Cosmochim. Acta* 74, 3571–3586.
- Zhao, Z.J., 1974. Red deer's living habits and its capture method. *Chin. J. Zool.* 9 (1), 22–23 (In Chinese).